



OVER EXPRESION OF THE *Swo1* GENE IN *Trichoderma atroviride* and EVALUATION OF ITS MICOPARASITIC ACTIVITY.

¹Karina Atriztán-Hernández, ¹Edgar Bálcazar-López, ²Alfredo Herrera-Estrella, ¹Jorge Luis Folch-Mallol; ²Laboratorio Nacional de Genómica para la Biodiversidad. Departamento de Ingeniería Genética de Plantas, CINVESTAV Irapuato. Apartado Postal 629, Irapuato 36821. ¹Universidad Autónoma del Estado de Morelos, Facultad de Ciencias Biológicas, Centro de Investigación en Biotecnología, Laboratorio de Biología Molecular de Hongos, Cuernavaca Morelos; C.P.62209; inatriztan@gmail.com

Key words: *Expansins*, *Swollenins*, *Trichoderma atroviride*.

Introduction. Expansins are proteins of plants, which are known to be involved in the remodeling of cell wall structures during growth and other processes (1). Other proteins with expansin-like activity have been identified in fungi, which have been named *Swollenins* (*Swo*), the first *Swo1*, was identified in *Trichoderma reesei* (2). Another *Trichoderma* species, *T. atroviride*, has been used as a model in phytopathogenic fungi control studies (3). A gene that encodes for a protein with similarity to *Swo1* was found to be highly expressed in the first stages in the process of mycoparasitism of *Trichoderma atroviride* with several phytopathogenic fungi (4).

Is of our interest to study the function of the *Swo1* gene of *T.atroviride* during mycoparasitism.

Methods.

The *Swo1* gene was obtained via RT-PCR from whole mycelium RNA of *T.atroviride* collected during direct confrontations with phytopathogens, later the gen was ligated to the TOPO-TA vector (Invitrogen) to transform to *Escherichia coli* DH5 α , and subcloned in the expression vector *pUE10*. Transformation of *T. atroviride* was made through a protoplasts technique; at least three monospores passes were performed to obtain stable transformants lines.

Results.




Fig.1 a) Confrontation of *T. atroviride* against a phytopathogenic fungus, b) *Swo1* gen amplified through RT-PCR, c) liberation of gen *Swo1* from the cloning vector TOPO-TA using restriction enzymes *NotI* y *EcoRI*.

The *Swo1* gene of *T. atroviride* was amplified and cloned (Fig1)



Fig. 2 a) Expression vector *pUE10*, b) gen *Swo1* liberation from expression vector with restriction enzymes *NotI* y *EcoRI*.

Figure 2(a) shows the expression vector *pUE10* used to subclone the *Swo1* gene in *E.coli* DH5 α , Figure 2(b) shows the verification of the *pUE10::Swo1* construction for the subsequent transformation into *T. atroviride*.

Conclusions:

The *Swo1* gene of cDNA of *T. atroviride* was amplified and cloned.

Sequence analysis showed that there are no nucleotide changes in *Swo1* clone.

The *Swo1* cDNA was introduced into the expression vector *pUE10* and with this construction *T. atroviride* was transformed by the protoplasts method.

Currently the analysis of the transformants strains is being performed.

Acknowledgements. This work was partially funded by grants CB 153789-Q, from CONACyT and SENER-CONACyT 150001

References.

1. Qiang Y., Ting-Ting S., Wei-Feng L., Guan-Jun C. (2008). *Biosci. Biotechnol. Biochem.* 72(11):2799-2805.
2. Saloheimo M., Paloheimo M., Hakola S., Pere J., Swanson B., Nyyssönen E., Bhatia A., Ward M., Penttilä M. (2002). *Eur. J. Biochem.* 269: 4202-4211.
3. Argumedo-Delira R., Alarcon A., Ferrera-Cerrato R., Peña-Cabriales J.J. (2009). *Rev. Int. Contam. Ambient.* 25(4):257-269.
4. Reithner B., Ibarra-Laclette E., L. Mach R., Herrera-Estrella A. (2011). *Appl. Environ. Microbiol.* 77(13):4361-4370.